# MagCore® Automated Nucleic Acid Extractor

# MagCore® Automated Nucleic Acid Extractor Overview

		tandard	Optional	Process Monitoring through your Smartphone  MagCore® Plus //	Spectrophotometer Built-in MagCore® Super	NA Extraction and PCR Set-up  Mag Core® EDA	
16 Sample	1-16Samples			-	-		
48 Sample	1-48 Samples					•	
Spectrophotometer	Spectrophotometer				-	-	
Touch Screen	Touch Screen			-	-		
	UV Decontamination				•	•	
Barcode Scanner	Barcode Scanner			-	•	-	
Thermo Printer	Thermo Printer				-		
LipgradiablevioLUS2	<b>Built-in Programs</b> (Upgradeablevia USB ports, Plug&Play)			-	-	-	
USBOutput	<b>USB Output</b> (USB flash drive not provided)					-	
Monitoring	<b>Progress Monitoring</b> (Wireless)						
UMS C	<b>LIMS</b> (Laboratory Information Management System)					-	
	HEPA Decontamination					•	
	Extraction +PCR set up						
	PCRsetup					•	
					!		

# MagCore® Automated Nucleic Acid Extractors will keep you ahead in Life Science Other Feature





MagCore® Extractor System is a simple, fast and cost-effective instrument for automated purification of nucleic acids from a diverse range of sample sources. Featuring pre-programmed protocols and our unique magnetic-bead technology, MagCore . System delivers efficient and consistent nucleic acid purification.

MagCore® Extractors are bench-top instruments ensuring efficient and cross-contamination free isolation of DNA/RNA. Built-in UV lamps allow to easily and efficiently decontaminate the instruments after run.

## **Flexibility**

MagCore® Automated Extraction System allows you to save time without sacrificing consistency and purity. You can use one instrument to purify DNA and RNA from a broad variety of sample types: from blood to mouse tails and almost everything in between.

### Ease Of Use

You will be provided with everything you need to run purifications, including pre-filled cartridges, specialized disposable tips and tubes. With the user-friendly interface and our user manuals, you are quaranteed to operate with ease.

MagCore® Automated Extraction System helps minimize cros-contamination by limiting hands-on procedures and turnaround time.

MagCore® System speeds the front-end processing, enabling you to do more tests in less time. And the Instrument is compact, soit can virtually fit into any lab.

# **Built-in Programs**

All of our MagCore Extractor models have built-in protocols for all of the kits we offer. Simply run the protocol by selecting the 3-digit codeprinted on the kit of interest.

Free upgrade of software and protocols can be downloaded from our website (www.rbcbioscience.com) and uploaded through the instrument RS232/USB ports.

## Diverse Sample Purification

We offer extraction kits designed for Blood, Plasma, Cell, Tissue, FFPETissue and Plant samples, to fit all your research needs.

Competitive Price and Small Footprint

# Easy To Use

Apply samples to instrument





Select the number of the cartridge.

D						
10	н	201				
10	)2	202				
▲ Prev Next		Cancel				
Select Cartridge Code						

Select Sample Volume



Push Start




A Beep Sound can be heard after protocol is completed



# Other Features

### Automatic Optical Density Measurement

- Built-in spectrophotometer provides O.D. A<sub>260</sub> and A<sub>280</sub> measurement of individual samples. (O.D. detection range: ABS < 6.)
- A<sub>320</sub> Normalization
- Disposable cuvettes.



### Thermo Printer and Barcode Scanner



### **Progress Monitoring**














MagCore® Nucleic Acid Extraction Kits

# MagCore® Automated Nucleic Acid Kits Selection Guide




# MagCore® Cartridge Design and Extraction Principle




MagCore® cartridges include all reagents needed for purifications, no additional handling is necessary. We minimize any possible contamination and spillage with an automated piercing step for our pre-sealed cartridges. RBC patented Heating Well and Separation Well in the cartridge provide a strong circular force to ensure efficient binding and washing during the extraction procedure.

MagCore® Nucleic Acid Extraction Kits contain all reagents and consumables needed for 36, 72 or 96 nucleic acid isolation reactions. The consumables consist of reagent cartridges, individually packaged tip sets, sample tubes and elution tubes.



# MagCore® Worldwide Patented Magnetic Beads

Our design: multiple core inside, cellulose coating.



Particlerange	~20µm (in water)
VSM	~40emu/g
Cellulosew%	~50%
Coresize	~150nm
CoreVSM	~80emu/g
Core material	Fe <sub>3</sub> O <sub>4</sub>
Bead Binding capacity	1 mg beads bind ~300µg Calfthymus DNA ~350µg human placenta DNA











RBC Bioscience Corp.
www.rbcbioscience.com
info@rbcbioscience.com

# MagCore® Automated Nucleic Acid Extractor

# Full traceability and mobile monitoring on your smartphone

# **MagCore®** Plus II



MagCore® Plus II is the newest robotic bench-top workstation for a fast and high-yield nucleic acid purification from virtually all molecular diagnostic, biological, clinical and forensic sample types. With small footprint, light weight, user friendly interface, and a broad range of entirely built-in programs with free upgrades, 1-16 samples can be isolated simultaneously at your fingertip. The instrument simplifies your daily routine providing full traceability of kits and samples, through real-time mobile monitoring and a complete report that can be downloaded on a computer at the end of each run.



# Worldwide Patented Magnetic Beads

Cellulose-coated magnetic beads, coupled with our patented binding and separation technology, guarantee high quality extracts



# Ideal for both DNA/RNA extraction

Built-in protocols are created for extracting nucleic acids from a wide range of samples, including whole blood, plasma (circulating free nucleic acid), tissue, bacteria, virus, plant and forensic.



# Throughput up to 16 samples per run

From cartridge piercing to final eluate, all steps are performed by the instrument, that allows running 1 to 16 samples at one time, for a time-saving and flexible performance.



# Full traceability of the samples and kits

A report in .csv format is generated at the end of each run and contains all relevant data: user's name, sample and kit barcode, protocol number, sample and elution volume, start and end time. The file, opened on a computer, can be subsequently processed by a LIMS.



# Real-Time Mobile Monitoring

During the run, the instrument HMI can be accessed via Wi-Fi from your smartphone/tablet through our App, to see real-time information about the run processing status, remaining time and errors. Android and iOS compatible.



# **WDecontamination**

The equipped UV lamp minimizes the risk of cross-contamination and ensures user and product safety.



# Built-in Programs (Upgradeablevia USB ports, Plug&Play)

MagCore® Plus II features built-in protocols for all the extraction kits we offer and is equipped with a USB port for free protocol and software upgrades.



# Barcode Scanner

For sample and kit tracking and monitoring and an easier organization of the test results.



Load Samples And Install Accessorie



Select User and Scan Barcodes:



Select the code of the cartridge.



Select Sample Volume And Eluate Volume



Program Cod
102

Dow Immedia Cotted NA

Date Internal Cotted NA

Date I

A Beep Sound can be heard when the program completes.



Open the run report on your computer







# Samethroughput, smaller size




### Barcode Scanner



Laboratory Information Management System (LIMS) Unidirectional LIMS device, Ethernet cable



# Cartridge Design and Extraction Principle



### Mobile Monitoring with Android and iOS App



# Specification

Model	Plus II
System Method	Cellulose coated magnetic beads
System Components	<ol> <li>Pipetting Unit: X and Y-axis movement for sample transfer and dispense.</li> <li>PLC module, HMI and Driver main board embedded in</li> <li>UV Light: power 8w, life duration 11,000 hrs</li> <li>Heating Block: RT-90°C</li> <li>Display Screen: 7-inch color touch panel</li> <li>Accessories: T-racks, cartridge racks, barcode scanner, waste box</li> </ol>
Power Supply	Voltage: AC 100V~240V; Frequency: 50/60Hz
Dimension	W523 x D602 x H605 (mm) / W21 x D23.7 x H23.8 (inches)
Net Weight	70kg/154.35lbs

# Operating Parameters

,	
Processing Capacity	1-16 samples per batch
Processing Time	30-90 minutes (depends on sample type and method)
Sample Volume	200 μl/400 μl/1,200 μl/ 4ml * depending on the program.
Elution Volume	30µl/40µl/60µl/100µl/150µl/200 µl * depending on the program.
Yield	Average 6µg Genomic DNA from 200µl human whole blood
Purity	DNA:O.D A <sub>260</sub> / <sub>280</sub> ratio 1.8 ± 0.1 RNA:O.D A <sub>260</sub> / <sub>280</sub> ratio 2.0 ± 0.2
Pipetting Accuracy	30-60µl 20%; 60-100µl 10%; 100-1000 µl 4%

# Operating Environment

5 , ,	1-16 samples per batch		15°C-35°C	
Processing Time	30-90 minutes (depends on sample type and method)	Temperatures allowed during		
Sample Volume	200 µl/400 µl/1,200 µl/ 4ml * depending on the program.			
	30µl/40µl/60µl/100µl/150µl/200 µl * depending on the program.	Temperatures allowed during operation		
Yield	Average 6µg Genomic DNA from 200µl human whole blood	Pollution Degree	level ?	
Purity	DNA:O.D A <sub>260</sub> / <sub>260</sub> ratio 1.8 ± 0.1 RNA:O.D A <sub>260</sub> / <sub>260</sub> ratio 2.0 ± 0.2		LCVC12	
Pipetting Accuracy	30-60µl 20%; 60-100µl 10%; 100-1000 µl 4%			



FDA (10055336) reaistered and CE-IVD certified (Instruments & Reagents)











Automatic Extraction and Smart PCR Setup

# MagCore® EDA



MagCore® EDA system is new generation instrument for nucleic acid extraction and PCR setup pre-treatment. MagCore® EDA can process up to 48 samples (in batches of 24), and a has a built-in spectrophotometer which makes it a complete solution to increase laboratory efficiency and simplify the workflow. Users can benefit from automatic purification, OD values retrieval, high NA concentration in final elute and compatibility with LIMS.

using MagCore extraction kits, MagCore® EDA offers consistent, reliable DNA or RNA extraction in 30–120 minutes, depending on the sample type.

# Worldwide Patented Magnetic Beads

Cellulose-coated magnetic beads, coupled with our patented binding and separation technology, guarantee high quality extracts.

## Ideal for both DNA/RNA extraction

Built-in protocols are created for extracting nucleic acids from a wide range of samples, including whole blood, plasma (circulating free nucleic acid), tissue, bacteria, virus, plant and forensic

# Automatic Optical Measurements of OD Values

Built-in spectrophotometer and our optical module provide users the option to automatically measure OD values and concentration offinal eluates upon completion of the nucleic acid extraction process.

## **UV** Decontamination

The equipped W lamp minimizes the risk of cross-contamination and ensures user and product safety.

# HEPA Decontamination

HEPA system is installed in the PCR set up zone to eliminate contamination and ensure the purity of the samples.

# Automatic Barcode Scanning


Barcode reading of primary tube simplifies the tracking of the samples and ensures traceability.

# Throughput up to 48 samples

From cartridge piercing to final elute, all steps are performed by the instrument, that allows running 1-48 samples. The instrument processed samples in batches of 24 at a time and has run setting for 24/48 sample to be processed at once, for time saving and flexible performance.

# Smart PCR setup

After pre-mix buffer configuration is set, the instrument automatically calculates the required volumes of the primers and perform liquid level detection. Pre-run report is generated to avoid errors and ensure optimal performance.







### Automatic Optical Density Measurement

- Built-in spectrophotometer provides O.D. A<sub>260</sub> and A<sub>280</sub> measurement of individual sa mples.
- (O.D. detection range: ABS < 6.)
- A<sub>320</sub> Normalization
- Disposable cuvettes.



### EDA System Connectivity

Ì

With LIMS compatibility EDA system can be connected to various thermocyclers and laboratory computer to set up automatic information exchange and analysis workflow.



Zone B provides automatic PCR set up which is compatible with different thermocylers for the further process. The system has 10 SBS positions, option of keeping premixes cool and the option of using 0.1 and 0.2 ml standard tubes, along from customised tubes from other PCR supplier.



Model	EDA V
System Method	Cellulose coated magnetic beads
System Components	<ol> <li>Pipetting Unit: dispensing, transferring, 2 X-Y axis model</li> <li>Electric Control: PLC model and Arm-based main board embedded in</li> <li>UV Light: power 8W, life duration 11,000hrs</li> <li>Contamination Prevention: HEPA filtration, Pipette tips with filter</li> <li>Heating Block: RT-95°C</li> <li>Nucleic Acid Concentration Detection Source: D2 lamp</li> <li>Detection Wavelength: 230nm, 260nm, 280nm</li> <li>Liquid detection sensor: PCR setup Zone</li> <li>Barcode scanning: Primary tube</li> <li>Accessories: T-rack, Cartridge racks, trash drawer, 96-well PCR plate rack, 32-well 1.5ml tube rack</li> </ol>
Power Supply	Voltage: AC 200-240V; Frequency: 50/60Hz
Dimension	W1215*D900*H860 (mm)
Net Weight	240 Kg / 529 lbs

# Operating Parameters

1-48 samples (in batches of 24)	
30-120minutes (depends on sample type)	
200/400/1200/4000 μl (application dependent)	
30/40/60/100/150/200 μl (application dependent)	
200ul whole blood (average 6ug gDNA) 400ul whole blood (average 12ug gDNA)	
DNA: O.D A <sub>360</sub> /A <sub>260</sub> ratio 1.8±0.1 RNA: O.D A <sub>360</sub> /A <sub>260</sub> ratio 2.0±0.2	
Extraction Zone: 40µl <5%; 60µl <2%; 100-900µl <1.5% PCR setup Zone: 5~20µl≤5%; 20~50µl≤1%	
	30-120minutes (depends on sample type) 200/400/1200/4000 µl (application dependent) 30/40/60/100/150/200 µl (application dependent) 200ul whole blood (average 6ug gDNA) 400ul whole blood (average 12ug gDNA) DNA: O.D A <sub>260</sub> /A <sub>280</sub> ratio 1.8±0.1 RNA: O.D A <sub>260</sub> /A <sub>280</sub> ratio 2.0±0.2  Extraction Zone: 40µl <5%; 60µl <2%; 100-900µl <1.5% PCR setup Zone:

# Operating Environment

1				
Processing Capacity	1-48 samples (in batches of 24)			
Processing Time	30-120minutes (depends on sample type)	Temperatures allowed during transportation, storage, and packaging		
Sample Volume	200/400/1200/4000 μl (application dependent)			
	30/40/60/100/150/200μl (application dependent)	Temperatures allowed during operation		
Yield	200ul whole blood (average 6ug gDNA) 400ul whole blood (average 12ug gDNA)	Pollution Degree	level 2	
	DNA: O.D A <sub>260</sub> /A <sub>260</sub> ratio 1.8±0.1 RNA: O.D A <sub>260</sub> /A <sub>260</sub> ratio 2.0±0.2	,		
	Extraction Zone: 40µl <5%; 60µl <2%; 100-900µl <1.5% PCR setup Zone: 5~20µl≤5%; 20~50µl ≤1%			











# MagCore® Automated Nucleic Acid Extractor

# The first intrument with built-in Spectrophotometer















Built-in spectrophotometer and our optical module provide users the option to automatically measure OD values and concentration of final eluates upon completion of the nucleic acid extraction process.

Test results can be saved in the instrument, downloaded through he USB port and/or printed by the thermal printer. USB Output (USB flash drive not provided)

USB Output allows users to conveniently save tests reports in excel format and upload system updates with a USB flash drive.

### Thermal Printer

Test reports are available in hard copy.

### Laboratory Information Management System (LIMS)

Test results are automatically saved after optical measurements. You can save up to 1,600 tests in LIMS. Data can be easily transferred to a printer or computer in the same network and the report file can be subsequently processed by a LIMS.



## Worldwide Patented Magnetic Beads

Cellulose-coated magnetic beads, coupled with our patented binding and separation technology, guarantee high quality extracts.



### Ideal for both DNA/RNA Extraction

Built-in protocols are created for extracting nucleic acids from whole blood, plasma, tissue cell, plant cell, bacteria cell and virus samples.



### **UV** Decontamination

The equipped UV lamp minimizes the risk of cross-contamination and ensures user and product safety.



## High Capacity of 16 Samples

The instrument and protocols allow running up to 16 samples at one time, providing time-saving and flexible operation.



### Built-in Programs (Upgradeable via USB ports, Plug&Play)

MagCore Super has built-in protocols for all of the extraction kits we offer. Simply run the protocol by selecting the 3-digit code printed on the kit of interest. MagCore® is equipped with a USB port. Free upgrade of software or protocols can be downloaded from our website (www.rbcbioscience. com).



## Touch Screen with User-Friendly Interface

An integrated 7-inch full-color touch screen with user-friendly interface offers ease in operation. Only one touch is required to run your daily work.



### Barcode Scanner

It enables sample tracking and monitoring throughout the entire purification process and helps organize test results.



## Progress Monitoring

Remote (wireless) HMI device automatically transfers the data to your Android smartphone.



### Load Samples



Install Accessories



Select the code of the cartridae.



Select Sample Volume



Press Start




A Beep Sound can be heard after the program is complete!











# Cartridge Design and Extraction Principle



# Automatic Optical Scanning and Retrieval

- The optical module provides O.D.  $A_{260}$  and  $A_{280}$  measurement of individual samples.(O.D. detection range: ABS < 6.)









Model	Super		\ /\				TYT	
System Method		d magnetic beads						
System Components	3. UV Light: pov 4. Heating Bloc 5. OD Detection 6. Detection So 7. Detection W 8. Display Scree	n Range: ABS 0 -2.5 burce: D2 lamp lavelength: 260nm, 280 en: 7-inch color touch p T-racks, cartridge racks	,000hrs Onm panel , cuvette racks, ba	rcode scanr	er, thern	nal printer		
Power Supply		0V~ 240V; Frequency: 50						
Dimension	W760 x D700 x I	H770 (mm) / W29.92 x	:D27.55 x H30.31 (	(inches)				
Net Weight	78kg / 171.99lbs							

# Operating Parameters

Processing Capacity	1-16 samples per batch	
Processing Time	30-90 minutes (depends on ample type and method)	
Sample Volume	200 µl/400 µl/1,200 µl /4ml * depending on the program.	-
Elution Volume	30µI/60µI/100µI/150µI/200 µI	
Yield	Average 6µg Genomic DNA from 200µl human whole blood	-
Purity	DNA:O.D $A_{260}/_{200}$ ratio 1.8 $\pm$ 0.1 RNA:O.D $A_{260}/_{200}$ ratio 2.0 $\pm$ 0.2	
Pipetting Accuracy	500µl≤4%	7

# Operating Environment

3 , ,	1-16 samples per batch			
Processing Time	30-90 minutes (depends on ample type and method)	Temperatures allowed during transportation, storage, and packaging		
	200 µl/400 µl/1,200 µl/4ml * depending on the program.	, , , ,		
Elution Volume	30µl/60µl/100µl/150µl/200 µl	Temperatures allowed during operation		
Yield Average 6µg Genomic DNA from 200µl human whole blood  Purity DNA: O.D A <sub>260</sub> / <sub>260</sub> ratio 1.8 ± 0.1 RNA: O.D A <sub>260</sub> / <sub>260</sub> ratio 2.0 ± 0.2				
		Pollution Degree		
	500µl≤4%			













# For selective DNA purification and size selection for NGS

MagCore® NGS-Clean up Kit is designed for fast DNA fragment cleanup and size selection. The cleanup program results in 80% or higher recovery while removing all adapter dimers in sample that is then applicable during the preparation of Next Generation Sequencing (NGS) library. For DNA fragment size selection, the program selectively purifies nucleic acid fragments from 100-200 bp or from 200-500 bp in just minutes which is also suitable for library preparation. The entire purification procedure is fully automated and can be performed in 35 minutes.

701

MagCore®NGS Clean-up Kit



# **Features**

- 1. Automated DNA Cleanup and Size Selection ~ 30min
- 2. Precise Size Range Selection
- 3. High Recovery Rate
- 4. Easy and Rapid Workflow
- 5. Consistent and Reliable

# **Applications**

The MagCore® NGS Clean up Kit is designed to purify DNA for various downstream applications, including:

- 1. DNA library peparation for NGS
- 2. PCRorreal-time PCR




# Performance

### Product Introduction and Average Recovery

Program	Program Description	Avarage Recovery (%)
701A	DNA Cleanup	>80%
701B	DNA Size Selection 200-500bp	40~60%
701C	DNA Size Selection 100-200bp	>50%

### Comparison of DNA Cleanup methods



M: 100bp Marker.


Lane 1, 4, 7: MagCore NGS-Clean up Kit clean up DNA sample with 701 A program.

Lane 2, 5, 8: Brand A manually cleanup DNA sample.

Lane 3, 6, 9: Brand V manually cleanup DNA sample.


MagCore NGS-Clean up Kit: results of the automatic cleanup of DNA sample are comparable with Brand A and Brand V manual kits, resulting in high purity product with >80% recovery.

# Human genomic library prepared with DNA Library Prep Kit was analyzed by an Agilent Technologies 2100 Bioanalyzer



Red Line: Before DNA Clean-up Blue Line: After DNA Clean-up

### Linear Increase in DNA yield



M: 100bp Marker.

Lane 1: MagCore NGS-Clean up Kit size select 200-500bp with 701B program.

Lane 2: Brand Amanually size's elect 200-500bp. Lane 3: Brand V manually size select 200-500bp.

MagCore NGS-Clean up Kit is completely aotomated and can effectively size select DNA from 200-500bp, while Brand A and Brand V contain larger DNA fragments in the final product.

# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/PlusII				MagCore® HF16 / Compact			
CatNo.		Cat No.	Di unning Timo	CatNo.		Cat No.	Dunania a Timo
36 preps		96preps	Running Time	36 preps		96preps	RunningTime

**701** 

Cartridge Code

### MagCore®NGS Clean-up Kit

For 50 µl sample volumes.

Contents: Pre-Filled Cartridges , Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 18 months

MNC-01	MNC-02	35 min	MNC-01	MNC-02	35 min









# MagCore® Nucleic Acid Extraction Kits



# For extraction of genomic DNA from human whole blood sample

MagCore® Genomic DNA Whole Blood Kit is designed for extraction of total DNA (including genomic, mitochondrial and viral DNA) from whole blood, plasma, serum and buffy coat. The pre-filled cartridge contains proteinase K, a chaotropic salt, and guanidine hydrochloride for cell lysis and protein degradation. They enhance the binding between cellulose-coated magnetic beads and DNA. High quality DNA is eluted by low salt elution buffer or water after the removal of contaminants. Purified DNA of approximately 20-30 kb is suitable for PCR or other downstream applications.

101

MagCore® Genomic DNA Whole Blood Kit (Speedy installation)



102

MagCore® Genomic DNA Whole Blood Kit



104

MagCore®Genomic DNA Large Volume Whole Blood Kit (1.2 ml)



106

MagCore® Genomic DNA Whole Blood Kit (For Genotyping)



### **Features**

- 1. High performance of purified DNA in downstream applications such as qPCR.
- 2. High analytical sensitivity
- 3. Cartridges are pre-filled and sealed to prevent contamination.
- 4. Walkaway processing improves work efficiency.

# **Applications**

The MagCore® Genomic DNA Whole Blood Kit is designed to allow automated processing of multiple sample types in the same run. Sample types include:

- 1. Fresh and frozen whole blood
- 2. Buffycoat
- 3. Bodyfluids


High quality DNA available for various downstream applications, including:

- 1. PCR and real time PCR
- 2. Genotyping or sequencing
- 3. SNP,STR




### DNA quality unaffected by anticoagulants

Anticoagulant	A <sub>260</sub> / <sub>280</sub>	Conc.(ng/μl)	
1. EDTA	1.87	99.3	
2. Sodium Citrate	1.87	86.1	
3. Lithium Heparin(4ml)	1.95	106.02 113.16	
4. Sodium Fluoride	1.91		
5. Lithium Heparin(9ml)	1.95	101.34	




### Consistent yield and purity



It is shown that there is a scalability of DNA isolation from blood samples (20, 50, 80, 100, 150, and 200 µl) by the MagCore® HF16 The amount of DNA was determined by  $A_{260}$  measurement.

# Kbps 10



# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/Pl	usll

Cat No. Running Time 96preps

### MagCore® HF16 / Compact / HF48

Cat No.	Cat No.	Di vonina Timo
36 preps	96preps	RunningTime

101

Cartridge Code

# MagCore®Genomic DNA Whole Blood Kit (Speedy installation)

For 200 and 400 µl sample volumes.

Contents: Pre-Filled Cartridges (Including Proteinase K), Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 12 months

Cat No.

36 preps

MGB400-01	MGB400-02	39 min (sample volume:200 μl) 50 min (sample volume:400 μl)	MGB400-01		MGB400-02	44 min (sample volume:200 μl) 57 min (sample volume:400 μl)
-----------	-----------	--	-----------	--	-----------	--

102

## MagCore® Genomic DNA Whole Blood Kit

For 200 and 400 µl sample volumes.

Contents: Pre-Filled Cartridges (Including Proteinase K), Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 18 months

39min (sample volume:200 µl) 44 min (sample volume:200 μl) MGB400-04 MGB400-03 MGB400-04 MGB400-03 50min (sample volume:400 µl) 55 min (sample volume:400 µl)

104

### MagCore®Genomic DNA Large Volume Whole Blood Kit (1.2 ml)

For 1200 µl sample volumes.

Contents: Pre-Filled Cartridges (Including Proteinase K), Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 18 months

MGB1200 76 min (sample volume:1200 µl) MGB1200 83 min (sample volume:1200 µl)

### MagCore® Genomic DNA Whole Blood Kit (For Genotyping)

MGB-400-08

For 200 and 400 µl sample volumes.
Contents: Pre-Filled Cartridges (Including Proteinase K), Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 18 months

MGB400-07

41 min (sample volume:200 µl) 53 min (sample volume:400 µl) \*optical detection is not provided

MGB400-07

MGB-400-08

44 min(sample volume:200 ul) 57 min(sample volume:400 µl) (Applicable models: HF16, Compact)

Product	Contents	Cat.No.
Proteinase K Set	11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
DN A	50 µl RNase A (50mg/ml)	RN050
nnaseA	130 ul RNase A (50 ma/ml)	RN130













# For extraction of free circulating DNA from human plasma or serum

MagCore® Plasma DNA Extraction Kit is designed for purifying free circulating DNA from human serum or plasma using MagCore® automated extraction systems. The kit contains all required reagents and labware for automated purification based on magnetic-particle technology. Reagents necessary for a complete process are supplied and pre-filled in the cartridges, which can be easily loaded into the MagCore®instrument.

105

MagCore® Plasma DNA Extraction Kit (1.2 ml)



# **Features**

- 1. High performance of purified DNA in downstream applications such as qPCR.
- 2. Efficient recovery of fragmented DNA
- 3. Pre-filled and sealed buffer cartridges prevent contamination.
- 4. No phenol or chloroform extraction
- 5. Efficient removal of contaminants and inhibitors.

# **Applications**

The MagCore® Plasma DNA Extraction Kit efficiently purifies free circulating DNA. Sample types include:

- 1. Human plasma
- 2. Human serum

High quality DNA available for various downstream applications, including:

- 1. PCR and real-time PCR
- 2. Next Generation Sequencing (NGS)




# Performance

## High-quality of free DNA from plasma and serum

Quality and quantity analysis of the free DNA by Agilent Bioanalyzer 2100. Superior quality DNA are available from plasma and serum samples by using MagCore® products.


Sample ID	Sample type	Target Conc. (pg/µl)	35 bp marker Conc. (pg/µl)	10380 bp marker Conc. (рд/µl)
1	Human plasma	77.94	125	<i>7</i> 5
2	Human serum	138.48	125	<i>75</i>



### High stability of free DNA

Quality and quantity analysis of the free DNA by Agilent 2100 Bioanalyzer. It is shown by the evidence that free DNA of high quality and stability is possible to extract from 5 different plasma samples by using MagCore\*procudts.

SampleID	Sample type	Target Conc.(pg/μl)	35 bp marker onc. (pg/μl)	10380 bp marker Conc. (pg/µl)
1	Human plasma	65.95	125	<i>7</i> 5
2	Human plasma	<i>76.63</i>	125	<i>7</i> 5
3	Human plasma	87.22	125	<i>7</i> 5
4	Human plasma	57.96	125	<i>7</i> 5
5	Human plasma	60.53	125	<i>7</i> 5



# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/PlusII				MagCore® HF16 / Compact / HF48			
Cartridas Codo		CatNo.	Running Time			Cat No.	Di vanina Tima
CartridgeCode		96preps				96preps	RunningTime

105 MagCore® Plasma DNA Extraction Kit (1.2 ml)
For 1200ul sample volumes.

For 1200µl sample volumes.
Contents: Pre-Filled Cartridges, Proteinase K, PK Storage Buffer, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

	MPD1200	74 min (sample volume:1200 µl) *optical detection is not provided	MPD1200	70 min (samplevolume:1200 µl)
1		opticuracteedornsriot province	 1	

# Enzyme Selection Guide

	Product	Contents	Cat. No.
	Proteinase K Set	11 mg Proteinase K,1.25 ml PK Storage Buffer	PK011
DN/ A	50µl RNaseA (50mg/ml)	RN050	
	130 ul RNase A (50mg/m))	RN130	









RBC Bioscience Corp. www.rbcbioscience.com info@rbcbioscience.com





# For genomic DNA extraction from cultured cells and amniotic fluid

MagCore® Cultured Cells DNA Kit is designed to extract genomic DNA from up to 5x10<sup>6</sup> cultured cells using MagCore® automated extraction systems. The kit contains all required reagents and labware for automated extraction using magnetic-particle technology. Reagents are supplied in prefilled cartridges, which can be easily loaded into the MagCore® instrument.

110

### MagCore® Cultured cells DNA Kit

Magnetic beads
Separation Well 2
Separation Well 1
Separation Well 2
Heatblock Well 2
Heatblock Well 1


### *l-eatures*

- 1. High performance of purified DNA in downstream applications such as gPCR.
- 2. Efficient isolation of DNA from up to 5 x 106 cells.
- 3. Cartridges are pre-filled and sealed to prevent contamination.
- 4. No phenol or chloroform extraction.
- 5. Efficient removal of contaminants and inhibitors.

# **Applications**

High quality DNA available for various downstream applications, including:

- 1. PCR and real-time PCR
- 2. Next Generation Sequencing (NGS)






## DNA quality analysis by Spectrophotometer

DNA of three different cultured cells were extracted by the employment of MagCore® HF16 and MagCore® Cultured Cells DNA Kit (200µl sample volume). The following data show the yield and purity of DNA of three samples.

Sample 1:COS7 (1.2x10<sup>7</sup>) Sample 2:Junkat (1x10<sup>6</sup>) Sample 3:HEK293 (1x10<sup>6</sup>)

	1	2	3
Yield (mg)	16.44	13.08	24.42
DNA purity (A <sub>260</sub> / <sub>280</sub> )	1.93	1.87	1.92



### Amniotic fluid DNA extration

### Result

DNA quality is checked by the NanoDrop spectrophotometer and agarose gel electrophoresis after DNA purification from amniotic-fluid cells. Cells were harvested from 10~15 ml amniotic fluid samples at pregnancy weeks 16-18 by centrifugation for 10 minutes at 3000 rpm.

Sample ID	Conc. (pg/µl)	A <sub>260</sub> / <sub>280</sub>	A <sub>260</sub> / <sub>230</sub>	Yield (µg)
(1) GPT2967P	22.68	1.84	1.07	1.36
(2) GPT2952P	<i>33.9</i> 8	1.92	1.57	2.04
(3) NP679P	33.65	1.80	1.34	2.02
(4) NP777P	39.22	1.86	1.20	2.35
(5) MP795P	22.05	1.79	0.97	1.32

by NanoDrop ND-1000

Table 1.DNA quality analysis by NanoDrop.

Sample ID (1) to (5) are different amniotic fluid samples. It is shown that MagCore® HF16 system can purify 1~2 µg DNA from 10~15 ml amniotic fluid samples and DNA purity is with the A<sub>260</sub>/<sub>280</sub> ratio of around 1.8±0.1




Figure 1. Genomic DNA isolations were run by gel electrophoresis on 1% agarose gel. Lane M: RBC 1kb ladder marker

Lane 1:GPT2967P

Lane 2: GPT 2952P

Lane 3: NP679P

Lane 4: NP777P

Lane 5: NP795P

# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/PlusII						MagC	.ore®HF16/C	ompact/HF48
Cartridge Code	CatNo.		Cat No.	Running Time	CatNo.		CatNo.	Runnina Time
CannageCode	36 preps		96ргерѕ		36 preps		96preps	nurilling little

110

### MagCore® Cultured Cells DNA Kit

For 200µl sample volumes (up to 5 x 10<sup>6</sup> cells)

Contents: Pre-Filled Cartridges, Proteinase K, PK Storage Buffer, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 18 months

MCC-01SP

MCC-02SP

MCC-02SP

MCC-02SP

MCC-02SP

MCC-01SP

MCC-01SP

MCC-01

MCC-02

Product	Contents	Cat. No.
Proteinase K Set	11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
RNase A	50 µl RNase A (50mg/ml)	RN050
RNaseA	130µl RNase A (50mg/ml)	RN130









# MagCore® Nucleic Acid Extraction Kits



# For extraction of cf DNA from 4ml plasma or serum

MagCore® Circulating DNA large volume kit is designed for purifying free circulating DNA from human serum or plasma using MagCore® automated extraction systems. The kit contains all required reagents and labware for automated purification based on magnetic-particle technology. Reagents necessary for a complete process are supplied and pre-filled in the cartridges, which can be easily loaded into the MagCore®instrument.

115

MagCore® Circulating DNA large volume kit (4ml)

Magnetic beads separation Well 2

Magnetic beads separation Well 1

Binding Buffer 1800µl

Binding Buffer 1800µl

Binding Buffer 1800µl

Beads Nixtule 1800µl

Wash 18uffer 1000µl

Wash 2 Buffer 1000µl

ddH,O 1000µl

Lysis Buffer 1000µl

Lysis Buffer 1000µl

Heat block Well 1

## *Features*

- 1. High performance of purified DNA in downstream applications such as aPCR.
- 2. Efficient recovery of fragmented DNA
- 3. Pre-filled and sealed buffer cartridges prevent contamination.
- 4. No phenol or chloroform extraction
- 5. Efficient removal of contaminants and inhibitors.

# **Applications**

The MagCore® Plasma DNA Extraction Kit efficiently purifies free circulating DNA. Sample types include:

- 1. Human plasma
- 2. Human serum

High quality DNA available for various downstream applications, including:

- 1. PCR and real-time PCR
- 2. Next Generation Sequencing (NGS)



# **Performance**

### High stability of free DNA

Quality and quantity analysis of the free DNA by Agilent 2100 Bioanalyzer. It is shown by the evidence that free DNA of high quality and stability is possible to extract from 5 different healthy people plasma samples by using MagCore products.

Sample ID	Sample type	Plasma (ml)	Qubit (ng/μl)	Ctvalue (GAPDH gene)	Bioanalyzer Conc. (pg/µl)	35 bp marker Conc. (pg/µl)	10380 bp marker Conc. (pg/µl)
1	Human plasma	4	0.262	18.52	127.21	125	<i>7</i> 5
2	Human plasma	4	0.250	18.59	124.62	125	75
3	Human plasma	4	0.220	18.78	111.11	125	75
4	Human plasma	4	0.290	18.45	153.24	125	75
5	Human plasma	4	0.278	18.49	130.97	125	<i>7</i> 5


# Sample 1 Sample 2

Quality and quantity analysis of the free DNA by Agilent Bioanalyzer 2100. Superior quality DNA are available from plasma samples by using MagCore products.


Comparison of free DNA from 1.2 ml and 4 ml plasma

SampleID	Kits Code	Sample type	Plasma (ml)	Qubit (ng/μl)	Ct value (GAPDH gene)	Bioanalyzer (pg/μl)	35 bp marker Conc. (pg/μl)	10380 bp marker Conc. (pg/µl)
1	105	Human plasma	1.2	0.061	23.86	24.70	125	<i>7</i> 5
2	105	Human plasma	1.2	0.058	23.96	22.98	125	<i>7</i> 5
3	115	Human plasma	4	0.203	20.04	117.0	125	<i>7</i> 5
4	115	Human plasma	4	0.194	20.26	104.7	125	<i>7</i> 5


Code 105:MagCore® Plasma DNA Extraction Kit Code 115:MagCore® Circulating DNA large volume kit









Sample 5




Sample 1





Sample 4



# MagCore® Automated Nucleic Acid Kits Specification

		MagCore®Super/	HF16Plus/PlusII	MagCore® HF16 / Compact		
CartridasCodo	Cat No.	CatNo.	Running Time Cat No.  24preps		Cat No.	Dunaina Tima
Cartridge Code	24 preps	96preps		24 preps		96preps

MagCore® Circulating DNA large volume kit

For 4000ul sample volumes

Contents: Pre-Filled Cartridges, Proteinase K, PK Storage Buffer, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 18 months

146 min (sample volume:4000 µl) MPD4000-01 MPD4000-03 146 min (sample volume: 4000 µl)

	Product	Contents	Cat.No.
	Proteinase K Set	11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
DN/ A	50µlRNaseA(50mg/ml)	RN050	
	130 ul RNase A (50mg/m))	RN130	









# MagCore® Nucleic Acid Extraction Kits



# For extraction of viral DNA/RNA from serum, plasma and cell-free body fluids

The MagCore® Viral Nucleic Acid Extraction Kit is designed for purifying viral DNA and RNA from serum, plasma, and cell free body fluids. MagCore® magnetic particle technology delivers high-quality DNA/RNA that is suitable for direct use in downstream applications such as amplification or other enzymatic reactions. To minimize the risk of cross-contamination, plastic consumables inside the Kit are DNase/RNase treated and the operation system is designed to individually process samples at the same time. Multiple protocols are installed on the instrument and optimized for different sample volumes.

202

MagCore® Viral Nucleic Acid Extraction Kit (Low PCR Inhibition)



210

MagCore® Viral Nucleic Acid Large Volume Extraction Kit (2.4 ml)



211

MagCore® Viral Nucleic Acid Large Volume Extraction Kit (1.2 ml)



## **Features**

- 1. High performance of purified DNA in downstream applications such as qPCR.
- 2. High analytical sensitivity.
- 3. Cartridges are pre-filled and sealed to prevent contamination.
- 4. Walkaway processing improves work efficiency..

# **Applications**

MagCore® Viral Nucleic Acid Extraction Kit is designed to allow automated processing of multiple sample types in the same run. Sample types include:


- 1. Cell-free body fluids
- 2. Plasma and serum
- 3. CSF
- 4. Urine



# Performance

### HBV detection by Real-Time PCR

The isolations of HBV nucleic acid from samples containing different concentrations of HBV were subsequently detected by Real-Time PCR



### Sensitivity test for HBV detection

	Lane 1	1x10°HBV serums.
	Lane 2	1x10 <sup>1</sup> HBV serums.
	Lane 3	1x10²HBV serums.
	Lane 4	1x10³HBV serums.
	Lane N	Negative control.
-	Lane M	RBC 100 bp DNA Ladder Marker.




Figure 1. Nested PCR results of HBV at different concentrations. Viral nucleic acids were purified from samples containing different amount of HBV using MagCore® Viral Nucleic Acid Extraction Kit.

## Sensitivity test for HCV detection

Lane 1	5x10°HCV serums.
Lane 2	5x10 <sup>1</sup> HCV serums.
Lane 3	5x10 <sup>2</sup> HCV serums.
Lane 4	5x10³HCV serums.
Lane 5	5x10⁴HCV serums.
Lane 6	5x10 <sup>5</sup> HCV serums.
Lane P	Positive control.
Lane N	Negative control.
LaneM	RBC 100 bp DNA Ladder Marker.



Figure 2. Nested PCR results of HCV at different concentrations. Viral nucleic acids were purified from samples containing different amount of HCV using MagCore® Viral Nucleic Acid Extraction Kit.

# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/PlusII

MagCore® HF16 / Compact / HF48

CatNo. CatNo. CatNo. Running Time

Cat No. Cat No. Running Time

202

### MagCore® Viral Nucleic Acid Extraction Kit (Low PCR Inhibition)

For 200 and 400 µl sample volumes.

Contents: Pre-Filled Cartridges, Proteinase K, PK Storage Buffer, Carrier RNA, Rnase Free Water, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes
Shalf life: 18 months

Shelf life: 18 months

MVN400-03 MVN-400-04 73 min (sample volume:400 μl)

62 min (sample volume:200 µl)
73 min (sample volume:400 µl)
MVN400-03
MVN400-04
57 min (sample volume:400 µl)
66 min(sample volume:400 µl)

210

### MagCore® Viral Nucleic Acid Large Volume Extraction Kit (2.4 ml)

For 2400 µl sample volumes.

Contents: Pre-Filled Cartridges, Proteinase K, PK Storage Buffer, Carrier RNA, Rnase Free Water, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

(special T-rack required for 5 ml sample tubes)

Shelf life: 18 months

MVN2400 90

90 min (sample volume:2400 µl)

211

### MagCore® Viral Nucleic Acid Large Volume Extraction Kit (1.2 ml)

For 1200 µl sample volumes.

Contents: Pre-Filled Cartridges, Proteinase K, PK Storage Buffer, Carrier RNA, Rnase Free Water, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

Shelf life: 18 months

MVN1200

80 min (sample volume:1200 µl)
\* optical detection is not provided

MVN1200

73 min (sample volume:1200 µl)

Product	Contents	Cat.No.
Proteinase K Set	11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
Carrier RNA Set	1 mg Carrier RNA, RNase Free Water	CR001













# For Extraction of Viral DNA/RNA From Serum, Plasma, Swabs and Cell-Free Body Fluids by Using MagCore® System

MagCore® Viral Nucleic Acid Kits are designed to isolate viral DNA and/or RNA from plasma, serum, swabs and cell-free body fluids. All plastic consumables included in the kits are DNase/RNase-free to minimize the risk of cross contamination. Our patented magnetic beads will bind to the short DNA fragments with high affinity ensuring that all important genetic information is collected efficiently making your daily isolation routine easier and faster. This kit also features Internal Control selection in its protocol, being compatible with any viral detection kit.

203

MagCore® Viral Nucleic Acid Extraction Kit (High Sensitivity)



## *Features*

- 1. Higher analytical sensitivity.
- 2. Consistent and reproducible results.
- 3. High performance of purified DNA/RNA in downstream applications such as qPCR.
- 4. Cartridges are pre-filled and sealed to prevent contamination.
- 5. Provides the option of Internal Control selection.


# **Applications**

MagCore® Viral Nucleic Acid Extraction Kit allows automated processing of multiple sample types in the same run, including:

- 1. Cell-Free Body Fluids
- 2. Plasma and Serum
- 3. CSF
- 4. Urine
- 5. Swabs




### Viral Nucleic Acid Extraction Performance Analysis



MagCore\*203 shows better performance on real-time PCR with approximately 1~2CtValueless ( $n \ge 3$ )

# Comparison of real-time PCR results of nucleic acid extraction by MagCore®, Brand Q and Brand R



Purified viral nucleic acid by MagCore® is ideal for the downstream real-time PCR. Viral samples processed by MagCore®

### Internal Control (IC) Selection

Mo	Sample Concentration (IU/mL) – HBV Signal / ICSignal					- NTC	
No.	Kit	Signal	5x10 <sup>6</sup>	5x10⁵	5x10⁴	5x10³	IVIC
1		HBV	22.06	25.39	28.58	30.53	
	MacCara® 202	+IC	29.99	31.54	31.17	31.55	
2	MagCore® 203	HBV	23.82	26.51	29.78	33.17	<i>-</i>
		+IC	31.19	30.79	30.51	32.73	

MagCore 203 program provides internal control (IC) selection, and the real-time PCR analyzed by CE IVD HBV Quantification kit shows IC signal within criteria (Ct value 30±3). This confirms success in MagCore 203 viral extraction step and low co-purification of PCR inhibitors that may cause false amplification patterns.

# MagCore® Automated Nucleic Acid Kits Specification

		MagCore®Super/H	F16Plus/PlusII		MagC	ore® HF16/C	Compact/HF48
Cartridge Code	CatNo.	Cat No.	Dunning Time	Cat No.		Cat No.	Dunning Time
CannageCode	36 preps	96preps	Running Time	36 preps		96preps	Running Time

203

### MagCore® Viral Nucleic Acid Extraction Kit (High Sensitivity)

For 200 and 400 µl sample volumes.

Contents: Pre-filled Cartridges, Proteinase K, PK Storage Buffer, Carrier RNA, RNase Free Water, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes. Shelf life: 12 months

MVN400-05		MVN400-06	57 Mins/With IC Selection: 58 Mins (sample volume:200 µl) 66 Mins/With IC Selection: 67 Mins (sample volume:400 µl)	MVN400-05		MVN400-06	56 Mins/With IC Selection: 57 Mins (sample volume:200 μl) 72 Mins/With IC Selection: 73Mins (sample volume:400 μl)
-----------	--	-----------	--	-----------	--	-----------	---

Product	Contents	Cat.No.
Proteinase K Set	11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
Carrier RNA Set	1mg Carrier RNA, RNase Free Water	CR001













# For extraction of plant DNA from plant tissues

MagCore® Genomic DNA Plant Kit is designed to extract DNA from plant tissues and cells using MagCore® automated extraction systems. The provided Filter Column Set can filtrate hard tissue samples to prevent tissue residues from dogging the pipette syringe. The kit contains all required reagents and labware for automated extraction using magnetic-particle technology. Reagents are supplied and pre-filled in cartridges, which can be easily loaded into the MagCore® instrument.

301

## MagCore® Genomic DNA Plant Kit



## Features

- 1. Lysis buffer can degrade large amounts of polysaccharides present on plant cell walls
- 2. Cartridges are pre-filled and sealed to prevent contamination.
- 3. No phenol or chloroform extraction and alchol precipitation.
- 4. Efficient removal of contaminants and inhibitors.

# **Applications**

The MagCore® Genomic DNA Plant Kit enables DNA purification from various samples types, including:

- 1. Plant cells
- 2. Plant tissues

High quality DNA available for downstream applications, including:

- 1. PCR and real-time PCR
- 2. Next Generation Sequencing (NGS)



## Yields of nucleic acid purified from various sources by MagCore® Genomic DNA Plant Kit

	R-/X/I					
Sample ID	Position	Yield (ng/μl)				
Citrus nobilisLour.	Leaf	36.5				
Carica papaya	Leaf	49.2				
Lycopersicon esculentum	Leaf	71.9				
Ctrullus lanatus	Leaf	23.4				
Anisogonium esculentum	Leaf	240.3				
Trachycarpus fortunei	Leaf	154.7				
Mentha piperita	Leaf	54.6				
Anthoceros punctatus	Leaf	38.9				
Calix babylonica	Leaf	80.7				
Prnus campanulata	Leaf	35.7				
Lxeris chinensis	Leaf	161.4				
Capsicum annuum	Leaf	13.9				
Vigna radiata	Seed	7.0				
Medicago sativa	Leaf	21.0				
Vigna angularis	Seed	5.8				
Agaricus bisporus	Whole	14.3				

# MagCore® Automated Nucleic Acid Kits Specification

		MagCore®Super/H		MagC	Core® HF16/C	Compact/HF48	
Cartridge Code	Cat No.	CatNo.	Running Time	Cat No.		Cat No.	Di vanina Tima
CarinageCode	36 preps	96preps		36 preps		96preps	Running Time

MagCore® Genomic DNA Plant Kit
For 50-100 mg fresh tissues or 5-20mg dried plant tissues
Contents: Pre-Filled Cartridges, Rnase A, GP1 Buffer, GP2 Buffer, Filter Column Sets, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

		Time: 33 min (sample volume:400 ul)			33min(samplevolume:400 ul)
MGP-01	MGP-02	*optical detection is not provided	MGP-01	MGP-02	(Applicable models : HF16, Compact)

Product	Contents	Cat. No.
Proteinase K Set	11 mg Proteinase K,1.25 ml PK Storage Buffer	PK011
RNase A	50µl RNaseA (50mg/ml)	RN050
nivaseA	130 ul RNase A (50mg/m))	RN130













# For extraction of genomic DNA from tissues, swabs, FFPE, stool, feed/soil and forensic samples

MagCore®Genomic DNA Tissue Kit is designed for purification of total DNA (including genomic, mitochondrial and viral DNA) from a variety of animal tissues or cells by using MagCore® auto-extraction instrument. The provided Filter Column can filtrate hard tissue sample or swab sample to prevent tissue residues to obstruct pipette tip during the process of MagCore®. The method uses pre-filled cartridges based on cellulose coated magnetic bead technology.

401

### 401 MagCore® Genomic DNA Tissue Kit



## Features

- 1. Suitable for a wide variety of sample types, including forensic samples
- 2. Consistent and reproducible results
- 3. Purified DNA of approximately 20-30 kb in length is suitable for PCR or other enzymatic reactions.

# **Applications**

MagCore® Genomic DNA Tissue kit allows processing multiple sample types in the same run, including:

- 1. Änimal Tissues: Solid animal tissue, Stool samples
- 2. Swabs
- 3. Feed and soil samples
- 4. Formalin-fixed, Paraffin-embedded (FFPE) tissues samples (manual pretreatment)
- 5. Cultured Yeast samples
- 6. Forensics samples



# Performance

### DNA yield gained from 10 mg of mice tissue samples and 0.25 cm tail

Mice tissue	Yield (ng/ul)
Liver	17.5
Spleen	17.5
Brain	10
Lung	12.5
Kidney	32.5
Tail	7.5

Table 1. Genomic DNA was purified from 10mg of mice tissue samples and 0.25cm tail.





Figure 1. Genomic DNA from 10mg of various animal tissues were extracted by the MagCore®401 Genomic DNA Tissue Kit. DNA was eluted in 60 µL and quantitated using Nanodropn-1000.






Figure 2. Agarose gel electrophoresis of the various animal tissue DNA was isolated from MagCore® automated extraction.; Lane M: RBC 1 kb DNA ladder Marker; Lane 1:Rat kidney;Lane 2:Rat liver;Lane 3-4:Pork liver;Lane 5-6:Fish tissue.

# MagCore® Automated Nucleic Acid Kits Specification

		MagCore®Super/H	Ма	gCore® HF16/C	Compact/HF48		
Cautidas Cada	CatNo.	Cat No.	D. wasain a Time	CatNo.	Cat No.	D. maria a Tiran	
Cartridge Code	36 preps	96preps	RunningTime	36 preps	96preps	Running Time	

401

### MagCore® Genomic DNA Tissue Kit

For tissue and forensic samples (up to 40mg)
Contents: Pre-Filled Cartridges, Filter Column Sets, GT Buffer, Proteinase K, PK Storage Buffer, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes Shelf life: 18 months

MGT-0	)1		MGT-02	33 min (sample volume:400 µl) * optical detection is not provided	MGT-01		MGT-02	33 min (sample volume:400 μl)
-------	----	--	--------	--	--------	--	--------	-------------------------------

# Enzyme Selection Guide

	Product	Cat. No.	
	Proteinase K Set	11 mg Proteinase K,1.25 ml PK Storage Buffer	PK011
RNase A		50µlRNaseA(50mg/ml)	RN050
		130 µI RNase A (50mg/m))	RN130







RBC Bioscience







# For Automated Extraction of Genomic DNA from Formalin-fixed Paraffin-embedded (FFPE) Tissue

MagCore® Genomic DNA FFPE One-Step Kit is designed to purify total DNA (including genomic, mitochondrial, and viral DNA) from Formalin-fixed Paraffin-embedded tissues via MagCore® Automated Extraction instruments. Our program features One-Step Heating, which automatically melts paraffin and lyses cells at the same time. No harmful reagent such as xylene is involved in the deparaffination process. Two protocols are outlined for different tissues sizes: 2-hour setup for smaller tissues and 16-hour setup for larger tissues.

405

405 MagCore® Genomic DNA FFPE One-Step Kit




## Features

- 1. Can load FFPE tissue samples with a surface area of up to 300 mm2 and up to 5 µm thick.
- A full package for DNA to be purified from FFPE samples. Complete automatic procedure from deparaffination to elution.
- 3. Stable deparaffination by Sula oil and reproducible results; no xylene involved.
- 4. Thermostavle cap is specially designed to cover the reaction well for an optimal condition where the lysis buffer and the FFPE sample can mix properly during the processing period.

# **Applications**

- 1. PCR and real-time quantitative PCR
- 2. Southern Blot
- 3. Next Generation Sequencing
- Genotyping





# Pertormance

# 2-hr heating protocol works out


There are 2-hour heating and 16-hour heating protocols. It is suggested for a better performance that the 2-hour heating program is for small tissues and the 16-hour program is for large


### Genomic DNA with high performance

The DNA extracted by MagCore® Genomic DNA FFPE One-Step Kit is high yield and suitable for downstream PCR and real-time PCR.

### A. DNA concentration


Whether samples were treated with either a 2-hour or 16-hour heating, yields of DNA extracted by MagCore® are mostly higher than the yields of DNA extracted by the competitor.






### B. The downstream PCR

PCR products of DNA purified by MagCore® and the competitor brand are confirmed by 1.5% agarose gel electrophoresis.









### DNA yield gained from 10 mg of mice tissue samples and 0.25 cm tail

Sample ID Sample ID	Yield (ng/ul)
Liver	17.5
Spleen	17.5
Brain	10
Lung	12.5
Kidney	32.5
Tail	7.5

### C. Purified DNA ideal for the downstream real-time PCR

Whether samples experience a 2-hour or 16-hour heating, the CT values of samples processed by MagCore® are mostly lower than the CT values of samples processed by the competitor.





It is shown below that different sizes of tissue samples were processed by a 2-hour or 16-hour heating program.

### 2-hourheating

### 16-hourheating

	Sample	Size(cm²)		Sample	Size(cm²)
1	319	1*1	1	128	<i>2</i> *3
2	317	0.8*0.8	2	191	2*2
3	170	0.5*1	3	350	2*2
4	171	1.5*1	4	362	<i>2</i> *3
5	354	1.5*2	5	369	<i>2</i> *2
6	349	1.5*2	6	304	<i>2</i> *3
7	207	0.5*0.5	7	206	1.5*1
8	208	0.5*1	8	207	0.5*0.5
9	209	0.5*1	9	208	0.5*1
			10	209	1*1

# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/Plus II MagCore® HF16 / Compact / HF48 Cat No. Cat No. Cat No. Cat No. Cartridge Code Running Time Running Time 36 preps 96ргерз 96preps 36 preps

### MagCore® Genomic DNA FFPE One-Step Kit

For FFPE tissue samples.
Contents: Thermostable Cap, Sula Oil, Pre-Filled Cartridges, Filter Column Sets, GT Buffer, Proteinase K, PK Storage Buffer, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes Shelf life: 18 months

175 min(2-hour heating)-Standard MGF-01 33min(samplevolume:400 ul) 1012 min(16-hourheating)-High Yield

	Product	Contents	Cat.No.
Proteinase K Set		11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
RNase A	50µlRNaseA(50mg/ml)	RN050	
	130 µl RNase A (50mg/ml)	RN130	









# MagCore® Nucleic Acid Extraction Kits



# For extraction of genomic DNA from forensic samples

MagCore® Genomic DNA Forensic Direct Kit is designed for purifying total DNA from forensic samples, such as dried blood spots, cigarette butts, cartilage, hair roots, seminal stains, and chorionic villus, using MagCore® auto-extraction instruments. Its unique feature is RBC patented technology that allows to isolate DNA automatically from solid samples without any pretreatment.


406

MagCore® Forensic DNA Direct Kit




# **Applications**

It uses magnetic-particle technology to purify genomic DNA from forensic samples. The purified genomic DNA can be directly used for downstream applications such as STR, PCR and real-time PCR.




# Automated Lysis and Separation





# **Performance** Unique Pipette Tip Design Airtight assurance (hermetic seal) Minimal aerosol contamination Automatic separation of the solid debris Screw Cap

## Compare the results between 406 and Qkit by qPCR



## Other casework samples

Reference samples





Figure: Extraction of simulated casework samples. This test was analyzed by real-time PCR with GAPDH gene primer. Saliva swab: 10 ml of saliva was applied (n=4). Blood swab: 1 ml of blood was applied and allowed to dry (n=4). Hair: 1 hair each from donor (n=4). Ggarette butt: A quarter of a filter paper (n=4).



Tipcollars

Designed hold

Top filter



# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/PlusII

Cat No. Cartridge Code Running Time 72preps

406

### MagCore®Forensic DNA Direct Kit

For extracting genomic DNA from forensic samples

Contents: Pre-Filled Cartridges (Including Proteinase K), Disposable Tip & Holder Sets, Elution Tubes, FS Tip, 200 ml SP Tip Shelf life: 12months

120 min

# Enzyme Selection Guide

Product	Contents	Cat.No.
Proteinase K Set	11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
DA1 4	50 µl RNase A (50mg/ml)	RN050
RNase A	130µl RNase A (50mg/ml)	RN130









RBC Bioscience Corp. www.rbcbioscience.com info@rbcbioscience.com





# For extraction of genomic DNA from bacteria

MagCore® Genomic DNA Bacterial Kit is designed to extract genomic DNA from Gram positive and Gram negative bacteria. The Kit contains all required reagents and labware for automated extraction using magnetic-particle technology. Reagents are supplied and prefilled in cartridges, which can be easily loaded into the MagCore® instrument.

502

# MagCore® Genomic DNA Bacterial kit

Magnetic beads
Separation Well 2

Magnetic beads
Separation Well 1

Empty
Lysis Buffer 1000µl

Beads Mixtu essay 1

Wash 1Buffer 1000µl

Wash 2Buffer 1000µl

Empty

Empty

Heat blockWell 1

Heat blockWell 1

## **Features**

- 1. High performance of purified DNA in downstream applications such as aPCR.
- 2. Cartridges are pre-filled and sealed to prevent contamination.
- 3. No phenol or chloroform involved.

# **Applications**


MagCore® Genomic DNA Bacterial Kit is designed for automated DNA extraction from various sample types, including:

- 1. Cell cultures in suspension (up to 5x10° cells)
- 2. Biological fluids
- 3. Cell cultures on plate
- 4. Gram-positive and Gram-negative bacteria




## High performance of purified bacterial DNA in real-time PCR

Well	Туре	Target	Ct	Interpretation	Label
D2	Unknow	H. pylori	30.70	Positive	Sample A
D3	Unknow	H. pylori	30.52	Positive	Sample A
D4	Unknow	H. pylori	NoCt	Negative	Negative Ctrl
D5	Unknow	H. pylori	30.59	Positive	Sample A
E2	Unknow	now H. pylori 26.73		Positive	Sample B
E3	Unknow	H. pylori	26.28	Positive	Sample B
E4	Unknow	H. pylori	26.40	Positive	Sample B



Consistent Yield, Purity, and Reproducibility in MagCore® Bacteria Genomic DNA Extraction





# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/Plus II					MagC	ore® HF16/C	Compact/HF48
Cartridge Code	Cat No.	CatNo.	Running Time	CatNo.		Cat No.	RunninaTime
	36 preps	96preps		36 preps		96ргерѕ	nuririiriy iii rie

502

### MaqCore®Genomic DNA Bacterial kit

For 1200µl sample volumes.

Contents: Pre-Filled Cartridges, Lysozyme Reaction Buffer, Proteinase K, PK Storage Buffer, Rnase A, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes

MBB-01		MBB-02	39 min (sample volume:200 µl) * optical detection is not provided	MBB-01	MBB-02	44min(samplevolume:200μl)
--------	--	--------	--	--------	--------	---------------------------

	Product	Contents	Cat. No.
Proteinase K Set		11 mg Proteinase K,1.25 ml PK Storage Buffer	PK011
DN/ A	50µl RNaseA (50mg/ml)	RN050	
	130 ul RNase A (50mg/m))	RN130	













# For extraction of microbial and host genomic DNA from stool samples

MagCore® Gut Microbiome DNA Kit is designed specifically for isolating high-quality microbial and host genomic DNA from stool and gut samples. With a short pretreatment, DNA from tough-to-lyse Gram-positive and Gram-negative bacteria which inhabit fecal samples can be successfully isolated with efficient lysis and homogenized with Beads-Beating Tube. In addition, inhibitor substances (e.g. polysaccharides, protein, etc.) commonly found in stools are removed with one-step IR Buffer. The crude DNA is then precipitated on ice and resuspended to adjust the condition that is suitable for binding genomic DNA.

Extracted DNA is ready for downstream molecular-based applications including qPCR, 16S rRNA gene sequencing, etc.

504

### MagCore® Gut Microbiome DNA Kit

Magnetic beads
Separation Well 1

Magnetic beads
Separation Well 1

Proteinase K 50µl

Empty
Binding Buffer GM
Buffer 1000µl

Wash 1 Buffer 1000µl

Wash 2 Buffer 1000µl

Empty
Empty
Empty
Empty
Empty
Heat block Well 1

# Features

- 1. Streamlined protocols for the isolation of inhibitor-free, PCR-quality DNA (up to 50 µg/prep) from microbes including Gram-positive and Gram-negative bacteria.
- 2. Hands-on time can be n as little as 20 minutes.
- 3. Depletion of PCR inhibitors by precipitation using a novel one-step IR Buffer.
- 4. High-purity DNA is ready for common downstream applications such as qPCR and 16S rRNA gene sequencing.

# **Applications**


The purified DNA can be directly used for downstream application such as:


- 1.aPCR
- 2. 16S rRNA gene Sequencing
- 3. Next-Generation Sequencing Applications



# Performance


### Stool DNA Extraction Efficiency






DNA was isolated from 0.2 g stool samples (in triplicates) with the MagCore Gut Microbiome DNA kit and supplier kits. (supplier T, supplier Z, supplier Q1, supplier Q2). DNA concentration and purity (A260/280, A260/230) are measured by Nanodrop spectrophotometer.

### **Gel electrophoresis**




Analysis of Genomic DNA extracted from human stool samples using 1% gel electrophoresis. Performance of MagCore Gut Microbiome kit analyzed on agarose gel electrophoresis; Lane M: RBC 1 kb DNA ladder Marker.

### 16SrRNA gene qPCR




DNA extracted from stool samples using MagCore 504 kit and other competitor kits were evaluated by 10-fold serial dilution of 16S rRNA gene qPCR amplification. Non template control (NTC) was included as a reference. (amplicon size: 466bp).



 $qPCR amplification curve of MagCore^*504, supplier Q2, supplier Z^*. MagCore 504 shows a low inhibition, similar amplification curve with supplier Q2, and superior Cq values to supplier Z.$ 

### Human fecal microbiota qPCR analysis



Phylum	Target organism	Gram Staining
Bacteroidetes	Bacteroides-Prevotella-Porphyromonas	G-
Actinobacteria	Bifidobacterium	G+
Actinobacteria	Clostridium coccoides–Eubacterium rectale group	G+
Firmicutes	Enterococcus spp.	G+

Human fecal microbiota qPCR analysis of four bacterial targets: Bacteroides-Prevotella-Porphyromonas, Bifidobacterium, Clostridium coccoides-Eubacterium rectale group, Enterococcus spp.. DNA from triplicate stool samples were isolated wih MagCore gut microbiome DNA kit and other suppliers.

# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/PlusII						MagC	Core® HF16/C	Compact/HF48
Cartridge Code	CatNo. Ca	atNo.		Running Time	Cat No.	Cat No.		Running Time
	24preps 72	2preps			-	-		nuririiriy iirrie

**504** 

### MagCore®GutMicrobiomeDNAKit

For 200±20 mg stool sample input.

Contents: Pre-filled cartridges, Beads-Beating Tube, GM-L1 Buffer, GM-L2 Buffer, GM-IR Buffer, GM-P1 Buffer, GM-P2 Buffer, GM-LY Buffer, Disposable tip & holder sets, Sample tubes, Elution tubes

Shelf life: 12 months











# For extraction of total RNA from formalin-fixed paraffinembedded (FFPE) tissue by using MagCore® System.

MagCore® Total RNA FFPE One-Step Kit is specially designed for total RNA purification from FFPE tissues by MagCore® instruments. It features the method, one-step heating, to melt paraffin without harmful reagents involved such as xylene or other organic solvents, and lyse tissues at the same time. The MagCore® Total RNA FFPE One-Step Kit System optimizes the lysis conditions to reverse the formalin fixation without the need for overnight digestion and retain both large and small RNAs. The program provides optional DNase I treatment to remove contaminated DNA.



### MagCore® Total RNA FFPE One-Step Kit



### Features


- A full package for total RNA isolation from FFPE tissue samples. It starts from melting paraffin to the final RNA purification.
- 2. Highly user-friendly protocol with minimal pretreatment required.
- 3. Sula oil (Deparaffinization solution): noxylene or other organic solvents involved.
- 4. Thermostable cap is especially designed to cover the reaction well for an optimal condition where lysis buffer and the FFPE samples can mix properly during the long processing period.
- 5. The total processing time is within approximately 160 mins.

# **Applications**

### The purified RNA can be directly used for downstream application such as:

- 1. cDNA synthesis
- 2. real-time PCR
- 3. RT-PCR
- 4. Microarray
- 5. Next-Generation Sequencing


### FFPERNA extraction efficiency




	Rat tissue	Average Yields (µg)	Average A <sub>260</sub> / <sub>280</sub>	Average A <sub>260</sub> / <sub>230</sub>
	Liver	3.57±1.38	1.96±0.11	1.26±0.16
ŧу	Kidney	2.25±0.62	1.82±0.12	1.03 ±0.15

Total RNA was purified from various FFPE rat tissues stored at -80°C for 3 months. RNA yield and quality from one (liver, kidney) 50 µm tissue section per sample was determined by Nanodropn-1000.

# MagCore® 605 competitors Tissue block A






		rissue block C								
	6 -					٦				
	5 -									
RNA Yields (µg)	4 -									
íeld	3 -									
RNA)	2 ·			_	_					
	1 .									
	0 -					┙				
		605	supplie	r Z supplier	Q supplier 1	Γ				
		1.99	2	1.91	2.13					

Ticcus block C


605 supplier Z supplier Q supplier Z supplier Q supplier Z supplier Q supplier Z supplier Q supplie


MagCore® 605 retains high RNA extraction efficiency in various rat liver tissue blocks compare to several FFPE RNA purification kit suppliers. The RNA yield and quality was quantified by Nanodrop n-1000.



# Pertormance

### RNA integrity






			Competitor Q		MagCore®605			
	Tissue Block	Sample	RNA yields (ng/µl)	RIN	Sample	RNA yields (ng/µl)	RIN	
ĺ	#1	Q1	97.37	2.2	R1	77.9	3.2	
	#2	Q2	<i>58.7</i>	2.2	R2	62.6	3.3	

### RNA integrity assessment

Total RNA isolation from rat liver 50µm tissue sections of FFPE tissue blocks were measured on Agilent 2100 bioanalyzer. Isolated RNA from MagCore® 605 kit maintains its integrity (18S, 28S peaks).


## MagCore® 605 competitors





Brand	Sample	nanodrop (ng/μl)	A <sub>260</sub> / <sub>280</sub>	A <sub>260</sub> / <sub>230</sub>	CtValue (GAPDH)
MagCore®	1	60.12	1.93	1.48	24.01
605	2	74.79	1.89	1.52	22.02
SupplierZ	3	31.21	1.96	1.49	26.61
Supplier Q	4	117.54	1.93	2.17	23.75
SupplierT	5	5.89	1.46	0.15	27.39


Comparison of total RNA extraction efficiency by using various FFPE RNA extraction kit suppliers. Total RNA was purified from one rat liver 50 µm tissue section of a single FFPE tissue block for each sample. The RNA yield and quality was quantified by Nanodropn-1000.

### RT-PCR performance











### Evaluation of RNA performance in RT-PCR

Five amplicons of different length (118, 206, 312, 400, 503, 613 bp amplicon size) of the gene Rpl4 performed using cDNA derived from FFPE rat liver samples purified by various suppliers.

# MagCore® Automated Nucleic Acid Kits Specification

MagCore®Super/HF16Plus/PlusII					MagCore® HF16 / Compact			
Cartridae Cada	Cat No.	Cat No.	Running Time	CatNo.	Cat No.	D		
Cartridge Code	24preps	72preps		24 preps	72preps	Running Time		



MagCore® Total RNA FFPE One-step Kit
For extraction of total RNA from formalin-fixed paraffin-embedded (FFPE) tissue by using MagCore® System.
Contents: Pre-Filled Cartridges, Disposable Tip & Holder Sets, Elution Tubes Shelf life: 12 months

		,			 
MRF-01	MRF-03	140min (without DNase I treatment) 158min (with DNase I treatment)	MRF-01	MRF-03	147min (without DNase I treatment) 165min(with DNase I treatment)

Product	Contents	Cat.No.
an is	For 36 reactions RNase-Free DNase I (Lyophillized): 1500 Kunitz units x 1 vial, 1 ml RNase-Free Waterx 1, 15 ml DNase I Reaction Buffe	DN036
DNase I Set	For 96 reactions RNase-Free DNase I (Lyophilized): 1500 Kunitz units x 2 vial, 1 ml RNase-Free Water x 2, 30 ml DNase I Reaction Buffer	DN096













# For purification of miRNA from serum and plasma

MagCore® Plasma miRNA Extraction kit is designed for purification of high purity microRNA from 400µl serum and plasma using MagCore® automated extraction instrument. All the kit components of plastic consumables are DNase/RNase-free pretreated, to eliminate the possibility of cross contamination.



### MagCore® Plasma miRNA Extraction Kit

Magnetic beads
Separation Well 1

Empty

Lysis Buffer 500,ul

Blanding Buffer 400,ul

Beads Mixture 1000,ul

Wash 5 Buffer 1000,ul

RNaseFreeMater 1000,ul

RNaseFreeMater 1000,ul

Empty

Empty

Empty

Heattblock Well 2

Heattblock Well 1

## **Features**

- 1. Purification of miRNA from 400 µl serum or plasma
- 2. RP Buffer to precipitate inhibitors
- 3. High performance of purified miRNA in downstream applications
- 4. Cartridges are pre-filled and sealed to prevent contamination


# **Applications**


Purified RNA ready to use in downstream applications such as:

- 1. Next-generation sequencing (NGS)
- 2. Real-time RT-PCR
- 3. Microarray analysis
- 4. Biomarker discovery


### High recovery of microRNA

Extraction results of microRNA isolation from 400 µl plasma from 3 different donors with an elution volume of 60 µl using MagCore Plasma miRNA Extraction kit (620) or Brand Q kit.





### **Detection of microRNAs**



micrioRNA was isolated from 400 µl plasma MagCore Plasma miRNA Extraction kit (620) or Brand Qkit.miR-150 and miR-16-1 was detected by Real-Time PCR.



# MagCore® Automated Nucleic Acid Kits Specification

MagCore® Super/HF16 Plus/Plus II						MagCore® HF16/Compact		
CautidaaCada	CatNo.	Cat No.		Running Time	CatNo.	CatNo.		D. venice Time
Cartridge Code	24 preps	72preps			24 preps	72preps		RunningTime



### MagCore® Plasma miRNA Extraction Kit

For 400 µl sample volumes Content: Pre-Filled Cartridges, RB Buffer, RP Buffer (10 ml), Disposable Pipet Tip plus Holder Sets, Sample Tubes, Elution Tube Shelf life: 18 month

MMR-01	NANAR-O3	40 min (without DN ase I treatment)	A A A A D O 1	MAAD OO	34 min (without DNase I treatment)
IVIIVIK-U I	IVIIVIK-U3	60 min (with DN/ara I traatmont)	IVIIVIK-U I	IVIIVIR-U3	57 min (with DNaco I troatment)















MagCore® triXact RNA Kit is specially designed for total RNA purification from the three most common sample types in diagnostics, research, and forensics: up to 1x10<sup>6</sup> cultured cells, a variety of tissues, or whole blood samples. The program provides optional DNase I treatment to remove residual DNA and extract high quality DNA-free RNA. Make use of its oustanding extraction performance, effiency, plus easy-to-follow protocols to simplify your every day extraction!



### MagCore®triXact RNA Kit

Empty
Lysis Buffer 1000µl
Beack Mixture 200µl
Beack Mixture 200µl
Beack Mixture 200µl
Wash 1 Buffer 1000µl
Wash 1 Buffer 1000µl
Wash 2 Buffer 1000µl
DEPCH<sub>2</sub>O 1000µl
Binding Buffer 1000µl
Wash 2 Buffer 1000µl
Heat block Well 1

## Features

- 1. Excellent RNA Yield and Purity shown in both  $A_{260}/A_{280}$  ratio and RIN value
- 2. Highly efficienct and user-friendly protocol with minimal pretreatment required
- 3. Capable of isolating RNA from 3 types of samples: Cells, Whole blood, Tissues
- 4. Optional DNase I treatment to remove residual DNA

# **Applications**

Purified RNA ready to use in downstream applications such as:

- 1. PCR
- 2. Real-time PCR
- 3. Microarray target preparation
- 4. Northern blotting
- 5. NGS analysis



### Total RNA from 1 x 10<sup>6</sup>Hela Cell Extract

No.	Kit	nanodrop ng/μl	260/280	260/230
1		207.15	2.08	1.86
2	Brand Q	182.85	2.07	1.78
3		174.90	2.11	1.63
4	Brand Z	171.52	2.03	2.24
5		210.45	2.02	2.22
6		164.76	2.05	2.17
7	MagCore*	264.69	2.04	2.11
8	triXact RNA Kit	250.61	2.02	2.17
9	1	237.44	2.03	2.17
10	Brand M	224.35	2.11	2.26
11		252.69	2.10	2.30
12		222.45	2.08	2.18

Table 1. Analysis of Cell RNA from MagCore® triXact Kit and others




Fig 1. Gel Electrophoresis of RNA from 1 x 106 Hela Cell Extract




Fig 3. Concentration comparison of RNA from 400 µl Human Whole Blood



Table 2. Analysis of Whole Blood RNA isolated by MagCore® triXact RNA Kit and others




Table 3. Extraction efficiency of RNA from Human Whole Blood with various anticoagulants

## Total RNA from 30 mg Fish Tissue




Fig 4. Concentration comparison of RNA from 30 mg Fish Tissue with/without DN as el treatment

# Specification

# MagCore® Automated Nucleic Acid Kits Specification

		MagC	Core®Super/HF16Plus/PlusII	MagCore® HF16 / Compact		
CartridaeCode	Cat No.	CatNo.	Physica Tipes	Cat No.	CatNo.	Dunning Time
Cartridge Code	24 preps	72preps	RunningTime	24 preps	72preps	RunningTime



### MaaCore®triXact RNA Kit

For total RNA extraction from cultured cells, human whole blood and animal tissues Contents: Pre-Filled Cartridges, RB Buffer, RBC Lysis Buffer, Filter column Set, Disposable Tip & Holder Sets, Sample Tubes, Elution Tubes Shelf life: 12 months

	MRX-01	MRX-03	48min (without DNase I treatment; starting volume: 400 µl) 81 min (with DNase I treatment; starting volume: 400 µl) * optical detection is not recommended	MRX-01	MRX-03	58min (withoutDNaseItreatment;startingvolume:400 µl) 73min (withDNaseItreatment;startingvolume:400 µl) (Applicable models:HF16, Compact)
--	--------	--------	--	--------	--------	--

Product	Contents	Cat.No.
Proteinase K Set	11 mg Proteinase K, 1.25 ml PK Storage Buffer	PK011
	For 36 reactions RNase-Free DNase I (Lyophillized): 1500 Kunitz units x 1 vial, 1 ml RNase-Free Water x 1, 15 ml DNase I Reaction Buffe	DN036
DNase I Set	For 96 reactions RNase-Free DNase I (Lyophilized): 1500 Kunitz units x 2 vial, 1 m I RNase-Free Water x 2, 30 m I DNase I Reaction Buffer	DN096









